

BENEFICIAL IMPACT OF PIGLETS VACCINATION WITH UNISTRAIN® PRRS BEFORE TRANSPORT TO FATTENING FARMS WITH HIGH RISK OF PRRS VIRUS INFECTION

Kim, J.H.¹; Kim, M.H.²; Seo*, S.W.²

¹Dodram Swine agricultural cooperative, Icheon-si. ²HIPRA Korea, Soengnam-si (Republic of KOREA)

*Corresponding author: sangwon.seo@hipra.com



The Reference
in Prevention
for Animal Health

INTRODUCTION

Porcine respiratory and reproductive syndrome virus (PRRSV) is an important pathogen causing huge economic impact in swine industry worldwide. Especially in the multisite system, PRRS can be a problem when fattening farms are located in high density pig producing areas with high risk of PRRSV infection¹. Through this study, the efficacy of UNISTRAIN® PRRS administered to 8 week of age piglets was evaluated in fattening farms with severe respiratory disorder caused by PRRSV.

MATERIALS AND METHODS

A 650-sow farm produced PRRS-free 30 kg piglets and transferred them to 2 fattening units that were surrounded by several swine farms. These neighbor farms were infected by type-2 PRRSV and problems of high mortality rate were observed. In this situation, piglet mortality was increased due to type 1 and 2 PRRSV infection. Accumulated mortality rate until slaughter was 18.1-24.3% and 14.4-15.3% in fattening farm A and B, respectively.

To solve this problem, a type-2 PRRS MLV (Ingelvac® PRRS) and a type-1 PRRS MLV (UNISTRAIN® PRRS, HIPRA) were evaluated by vaccinating different piglet batches at 8 weeks of age (600-750 piglet per batch). Cumulative mortality rate of each batch was recorded to evaluate the efficacy of PRRS vaccines in terms of clinical signs and productivity.

RESULTS

As a first prevention measure, type-2 PRRS MLV was used during 4 weeks, but pig mortality remained still high in farm A (13.3%, Table 1) and B (9.9%, Table 2). Moreover, field strains of type-1 PRRSV were detected in serum samples by RT-PCR.

In order to control PRRSV infection, vaccination with UNISTRAIN® PRRS was implemented. Using UNISTRAIN® PRRS, cumulative mortality was significantly reduced in farm A and B during 5 months. Especially after the 3rd batch of piglets vaccinated with UNISTRAIN® PRRS, cumulative mortality rate until slaughter age in farm A and B decreased until 2.3% and 2.1%, respectively.

Table 1. Cumulative mortality rate of each batch in farm A.

GROUP	BATCH SIZE (number of pigs)	ENTRY DATE (30 kg pigs)	CUMULATIVE MORTALITY RATE (until slaughter age)
Non-vaccinated	700	16 Jul 16	24.3%
	700	06 Aug 16	18.1%
Ingelvac® PRRS	750	13 Sep 16	13.3%
	750	25 Sep 16	8.5%
UNISTRAIN® PRRS	750	30 Oct 16	6.5%
	706	24 Nov 16	2.3%
	650	24 Dec 16	3.2%
	699	25 Jan 17	3.2%
	650	10 Feb 17	2.2%

Table 2. Cumulative mortality rate of each batch in farm B.

GROUP	BATCH SIZE (number of pigs)	ENTRY DATE (30 kg pigs)	CUMULATIVE MORTALITY RATE (until slaughter age)
Non-vaccinated	700	04 Jul 16	14.4%
	700	26 Jul 16	15.3%
Ingelvac® PRRS	750	18 Aug 16	9.9%
	750	01 Sep 16	7.6%
UNISTRAIN® PRRS	750	07 Oct 16	3.3%
	706	19 Oct 16	2.1%
	650	11 Nov 16	1.9%
	699	08 Dec 16	2.0%
	650	08 Jan 17	2.1%




Figure 1. Cumulative mortality rate in fattening farm until slaughter age (Gray; Non-vaccinated, Blue; Ingelvac® PRRS, Orange; UNISTRAIN® PRRS).

CONCLUSION

The commercial type-2 PRRS MLV could not control the type-1 PRRSV infection. However, in the same production system, UNISTRAIN® PRRS showed significant improvement in productivity reducing significantly the mortality. In overall, piglet vaccination with UNISTRAIN® PRRS improves productivity especially in farms co-infected with type 1 and 2 PRRSV².

REFERENCES

- Pauline Berton, Valérie Normand (2017). *Evaluation of porcine reproductive and respiratory syndrome stabilization protocols in 23 French Farrow-to-finish farms located in a high-density swine area.* Porcine Health Management 3:11
- Seongsik Ko, Sangwon Seo, *Efficacy of commercial genotype 1 porcine reproductive and respiratory syndrome virus (PRRSV) vaccine against field isolate of genotype 2 PRRSV.* Veterinary Immunology and Immunopathology 172:43-49